

PROVASTRIDE™ EFFICACY

EPA Reg. No. 63761-11

THE MOST COMPREHENSIVE CLAIM SET OF ANY REGISTERED, QUAT-FREE DRY SANITIZER*

Sanitization <i>5 minute contact time</i>		Disinfection <i>10 minute contact time</i>	Virucidal <i>90 second contact time</i>
<i>Klebsiella aerogenes</i>	<i>Escherichia coli</i>	<i>Pseudomonas aeruginosa</i>	Avian influenza A virus
<i>Staphylococcus aureus</i>	<i>Listeria monocytogenes</i>	<i>Staphylococcus aureus</i>	Feline calicivirus
<i>Salmonella enterica</i> <i>serovar Enteritidis</i>	<i>Salmonella enterica</i> <i>(Salmonella)</i>	<i>Salmonella enterica</i> <i>(Salmonella)</i>	Porcine Epidemic Diarrhea virus (PEDv)
<i>Cronobacter sakazakii</i>	<i>Campylobacter jejuni</i>		

*ProvaStride is approved as a dry, non-food contact sanitizer. ProvaStride can be applied as a registered sanitizer in its packaged form or can be pre-diluted in water and applied and a disinfectant to non-food contact surfaces.



Sterilex products are available from sanitation chemical providers. Please contact us or visit sterilex.com/presenting-provastride for more information.

111 Lake Front Drive, Hunt Valley, MD 21030 | 1.800.511.1659 | marketing@sterilex.com

The content in this marketing material is provided for informational purposes only and does not supersede the product label requirements. Approved label claims may vary depending on geography, use site, organism, or other factors. Always refer to the product label for complete directions for use.

PS_FE_120922